#### **INTERNATIONAL JOURNAL OF MEDICAL AND APPLIED SCIENCES**



E-ISSN:2320-3137

### **CASE REPORT**

# SUPERIOR MESENTRIC ARTERY SYNDROME [WILKIE'S SYNDROME]: A RARE CAUSE OF INTESTINAL OBSTRUCTION.

## G.V.Ramana Reddy\*, Siddartha Reddy, Abhilash Vemula, Sridhar Reddy, M, Pradeep Kumar.

Department of General Surgery, SVS MedicalCollege, Mahabubnagar, Telangana, India – 509 001

Corresponding author: Dr.G.V.Ramana Reddy, Dept of General Surgery, SVS Medical College and Hospital, Yenugonda, Mahabubnagar, 509001.

#### **ABSTRACT:**

Superior mesenteric artery syndrome is an entity generally caused by the loss of the intervening mesenteric fat pad, resulting in compression of the third portion of the duodenum by the superior mesenteric artery. It is a life-threatening upper gastrointestinal disorder due to compression of duodenum as it poses a difficult diagnostic dilemma. Third part of duodenum is in fixed compartment bounded anteriorly by the root of mesentery and superior mesentery artery and posteriorly by the aorta and lumbar spine.

**KEYWORDS:** Compression, third part of duodenum, Superior mesenteric artery, Aorta, Life threatening, diagnostic dilemma.

#### **INTRODUCTION**

Superior mesenteric artery syndrome (SMAS) is a rare cause of small bowel obstruction, characterized by an extrinsic vascular compression of the third portion of the duodenum between the abdominal aorta and overlying superior mesenteric artery, due to loss of cushion of fat. Clinically, it presents with postprandial abdominal distension, pain, nausea, vomiting, and weight loss. Fasting, total parenteral nutrition, and gastric decompression constitute usual conservative treatment with a high success rate. Surgery if needed has a low failure rate and consists of creating a gastrojejunostomy or duodenojejunostomy with or without duodenal mobilization (known as the Strong's procedure).

#### **CASE REPORT:**

A 17 year young man was admitted into the emergency with the complaint of recurrent abdominal pain, epigastric fullness, and vomiting and weight loss. Abdominal examination revealed upper abdomen fullness, resonant note on percussion and hyper peristaltic bowel sounds on auscultation. Erect abdomen X ray revealed dilated stomach bubble with non visualization of rest of bowel loops .CT scan of the abdomen revealed dilatation of stomach, first and second part of the duodenum proximal to the aorta-superior mesenteric artery angle. Laboratory

Volume 4, Issue 3, 2015

#### **INTERNATIONAL JOURNAL OF MEDICAL AND APPLIED SCIENCES**



E-ISSN:2320-3137

investigations were normal. Conservative management was tried for 15 days but to no avail. Explorative laprotomy performed revealed compression of 3<sup>rd</sup> part of duodenum. The duodenal obstruction was bypassed by performing duodenojejunostomy. Post op recovery was uneventful.

#### **DISCUSSION:**

The superior mesenteric artery (SMA) arises from the anterior surface of the abdominal aorta, just inferior to the origin of celiac trunk, and supplies the intestine from the lower part of the duodenum through two thirds of the transverse colon, as well as the pancreas.

Superior mesenteric artery syndrome is a rare cause of upper gastro intestinal obstruction [1].only 400 cases have been reported in medical literature [2].

Superior mesenteric artery syndrome was described for the first time by Rokitansky in 1842 and has an incidence of 0.013–0.3% in the general population with a mortality rate of 33% [3, 4]. Other names also used to indicate this clinical entity include Wilkie's syndrome [5], aortomesenteric artery compression [6], arteriomesenteric duodenal compression [7], duodenal vascular compression [8], and cast syndrome [8].

The SMA usually forms an angle of approximately  $45^{\circ}$  (range,  $38-56^{\circ}$ ) with the abdominal aorta, and the third part of the duodenum crosses caudal to the origin of the SMA, coursing between the SMA and aorta. Any factor that sharply narrows the aortomesenteric angle to approximately  $6-25^{\circ}$  can cause entrapment and compression of the third part of the duodenum as it passes between the SMA and aorta, resulting in SMA syndrome. There are several factors that can decrease the acuity of the angle between the aorta and SMA; the most common is significant weight loss leading to loss of the mesenteric fat pad. As a result, SMA syndrome has most often been described in patients with severe, debilitating illnesses, such as malignancy or malabsorption syndromes. It has also been described in a variety of other disorders associated with extreme weight loss including anorexia nervosa, trauma or burns, spinal cord injury and paraplegia, and after prolonged bed rest (9,10). Other causes include high insertion of the duodenum at the ligament of Treitz, a congenitally low origin of the SMA and compression of the duodenum caused by peritoneal adhesions, due to duodenal malrotation (11, 12).

Diagnosis depends on high index of suspicion since symptoms can be non-specific. The syndrome usually affects young females (10 to 39 years). The symptomatology is commonly chronic, with epigastric pain, bloating after meals, and vomiting.

The clinical diagnosis can be confirmed by radiologic studies in 95% of cases (13). Historically, barium meal and arteriography were used as diagnostic tools (14) but more recently, CT, CT-angiography and magnetic resonance imaging (MRI) have been used and shown higher diagnostic sensitivity (15).

Arteriographic criteria include a significantly decreased aorta-SMA angle of  $6^{\circ}$  to  $25^{\circ}$  (NL= $45^{\circ}$ ) and a shortened aortomesenteric distance of 2 to 8mm (NL=10 to 20mm) (16).

Conservative management is the rule for acute cases (17). Conservative treatment includes adequate nutrition, G.I. decompression and proper positioning after eating. Prokinetic drugs like metoclopramide or cisapride may be helpful. Surgery is indicated for chronic cases and failure of conservative management.

The most common operation for SMA syndrome, duodenojejunostomy, was first proposed in 1907 by Blood good (18). This open surgery involves the creation of an alternate route between

Volume 4, Issue 3, 2015



the duodenum and the jejunum, bypassing the compression caused by the AA and the SMA. Laparoscopic duodenojejunostomy for the management of SMA has also been described in the literature (19).

#### **CONCLUSION:**

Superior mesenteric artery syndrome (SMAS) is a rare condition caused by compression of the transverse portion of the duodenum between the superior mesenteric artery (SMA) and the aorta. Symptoms are non-specific and the diagnosis depends on high index of suspicion. MRA, being not invasive, is rapidly replacing the arteriogram in confirming the diagnosis. Conservative management may be sufficient in early cases. Duodenojejunostomy is the surgical treatment of choice.



DILATED STOMACH AND DUODENUM FIG 1



DUODENOJEJUNOSTOMY FIG 2

CT SCAN FIGURE 3

Dilated Stomach and Duodenum

Volume 4, Issue 3, 2015

#### **INTERNATIONAL JOURNAL OF MEDICAL AND APPLIED SCIENCES**



E-ISSN:2320-3137

#### **REFERENCES:**

- 1. Palanivelu C, Rangarajan M, Senthilkumar R, Parthasarathi R, Jani K. Laparoscopic duodenojejunostomy for superior mesenteric artery syndrome. J Soc Laparoendoscopic Surgeon 2006; 10: 531-4.
- Roy A, Gisel JJ, Roy V, Bouras EP. Superior mesenteric artery (Wilkie's) syndrome as a result of cardiac cachexia. J Gen Intern Med 2005; 20: 3-4.
- 3. C. Rokitansky, Handbuch der pathologischen Anatomie, vol. 3, Braunmüller & Seidel, Wien, Austria, 1st edition, 1842.
- Valdes, O. Cárdenas, A. Espinosa, O. Villazón, and V. Valdes, "Superior mesenteric artery syndrome," Journal of the American College of Surgeons, vol. 201, no. 5, p. 808, 2005
- 5. R. H. Guthrie, "Wilkie's syndrome," Annals of Surgery, vol. 173, pp. 290–293, 1971.
- 6. U. A. Dietz, E. S. Debus, L. Heuko-Valiati et al., "Superior mesenteric artery syndrome," Chirurg, vol. 71, no. 11, pp. 1345–1351, 2000.
- W. C. Anderson, R. Vivit, I. E. Kirsh, and H. B. Greenlee, "Arteriomesenteric duodenal compression syndrome. Its association with peptic ulcer," The American Journal of Surgery, vol. 125, no. 6, pp. 681– 689, 1973.
- G. Sapkas and J. P. O'Brien, "Vascular compression of the duodenum (cast syndrome) associated with the treatment of spinal deformities. A report of six cases," Archives of Orthopaedic and Traumatic Surgery, vol. 98, no. 1, pp. 7–11, 1981.
- Pentlow BD, Dent RG. Acute vascular compression of the duodenum in anorexia nervosa. Br J Surg 1981; 68: 665.
- 10. Laffont I, Bensmail D, Rech C, et al. Late superior mesenteric artery syndrome in paraplegia: case report and review. Spinal Cord 2002; 40: 88.
- 11. Wilkie DPD. Chronic duodenal ileus. Am J Med Sci 1927; 173: 643-649.
- 12. Ortiz C, Cleveland RH, Blickman JG, et al. Familial superior mesenteric artery syndrome. Pediatr Radiol 1990; 20: 588-589.
- 13. Hearn JB. Duodenal ileus with special reference to superior mesenteric artery compression. Radiology 1966; 86: 305.
- Goin LS, Wilk SP. Intermittent arteriomesenteric occlusion of the duodenum. Radiology 1958; 67: 729-733.
- 15. Mansberger AR Jr, Hearn JB, Byers RM, Fleisig N, Buxton RW. Vascular compression of the duodenum.
- 16. Baltazar U, Dunn J, Floresguerra C, Schmidt L, Browder W. Superior mesenteric artery syndrome: An uncommon cause of intestinal obstruction. South Med J 2000; 93: 606-608.
- 17. Palanivelu C, Rangarajan M, Senthilkumar R, Parthasarathi R, Jani K. Laparoscopic duodenojejunostomy for superior mesenteric artery syndrome. J Soc Laparoendoscopic Surgeons 2006; 10: 531-4.
- 18. Baltazar U, Dunn J, Floresguerra C, Schmidt L, Browder W. Superior mesenteric artery syndrome: an uncommon cause of intestinal obstruction. South Med J 2000; 93: 606-8. PMID 10881780.
- 19. Selami S, Hayrunisa K, Arzu A, Gulsen E, Haluk GB. Incomplete duodenal obstruction in a newborn. Indian J Paediatr 2006; 73: 364-6.