# **Original Research Article**

# COMPARATIVE EVALUATION OF ANTIFUNGAL ACTIVITY OF COCOS NUCIFERA OIL AGAINST CANDIDA ALBICANS

# Natarajan Kannan<sup>\* 1</sup>, Ajmal Mohammed<sup>2</sup>

1. Professor & Head, 2. Postgraduate Student

# Department Oral Medicine & Radiology, Narayana Dental College, Nellore, A.P., India.

#### Corresponding Author: Dr.Natarajan Kannan

#### **ABSTRACT:**

Candidiasis is a fairly frequent clinical finding in patients with xerostomia, radiotherapy patients, patients under steroid therapy and denture wearers. Edible coconut oil is a readily available substance readily available around the world with ability to reduce burning sensation in mouth and also some amount of lubrication in oral cavity. This study was carried out with the aim of assessing antifungal activity of virgin coconut oil and to determine its efficiency in treatment of oral candidiasis as a natural antifungal agent. Our study showed that virgin coconut oil exerts an antifungal effect on Candida albicans. The findings as reported in the present study indicate a strong potential therapeutic value of virgin coconut oil against Candidal infections.

Keywords: Virgin Coconut oil, Oral Candidiasis, Antifungal activity, Agar well diffusion technique

#### INTRODUCTION

Candida albicans is a dimorphic yeast that can be either a commensal or an opportunistic pathogen with the ability to cause a variety of infections, ranging from superficial to life threatening.<sup>1, 2</sup> C. albicans lives as a normal commensal in the oral cavity in 30 to 60% of healthy individuals.<sup>3</sup> C. albicans can cause infections under altered physiological and pathological conditions such as infancy, pregnancy, diabetes; prolonged broad spectrum antibiotic administration, steroidal chemotherapy as well as AIDS.<sup>4, 5</sup> The difficulty associated with the management of Candidial infections necessitates the discovery of new antifungal agents in order to widen the spectrum of activity against Candida.

Widespread use of limited antifungal agents to treat infections has led to the rapid development of drug-resistant strains. Another important reason for the low success rate in treating candidiasis is with cells in biofilms displaying characteristics dramatically different from their free-living planktonic counterparts. Of all the strategies which have been identified to overcome resistance to antifungal drugs the search of new and effective natural products showing antifungal activity against biofilm cells with low cytotoxicity.<sup>6,7,8</sup>

Coconut oil is an edible extract from the nut of Cocos Nucifera. In Ayurveda coconut oil is used internally & externally as part of several treatment procedures. The lauric acid content of coconut endows it with antimicrobial properties. As such, coconut

www.earthjournals.org

Volume 4 Issue 2, 2014

oil is used in the treatment of digestive tract infections. It is also used to expel intestinal parasites like tapeworms and Helicobacter pylori, which are responsible for indigestion and ulcer.<sup>9,10</sup>

#### AIM:

The main purpose of our study is to evaluate the in vitro antifungal activity of virgin coconut oil and to determine its efficiency in treatment of oral candidiasis as a natural antifungal agent.

## MATERIALS AND METHODS:

The antifungal activity was evaluated by the agar–well diffusion method. Candida albicans ATCC 66396 were sub-cultured onto Sabouraud Dextrose Agar and incubated for 18 hours at 37°C. The yeast cultures were harvested and then suspended in sterile saline (0.8% NaCl) and the cell density was adjusted to  $10^7$  cells/ml. Six millimeters – diameter wells were punched into the agar plate.

In Group A: 500  $\mu$ l of virgin coconut oil was dispensed into the wells. Same volume of clotrimazole (1%) was used on another plate as control (Group B). The plates were then incubated at 37°C for 24 hours. The diameters of the zones of inhibition around each disc were examined, measured and recorded as the mean diameter (mm) of complete growth inhibition. Tests were done in triplicate and the results are given as the mean average

#### **RESULTS:**

Zone of inhibition measured at 24 hours in test groups (A1, A2, and A3) with a mean average of 19.6mm [fig 1]. Zone of inhibition of the control group of clotrimazole (B1, B2 and B3) with a mean diameter of 23.2mm [fig 2]



## Culture Plates showing Agar Disc Diffusion Method

Zone of inhibition measured at 24 hours in test groups i.e. with Virgin Coconut Oil (A1, A2, and A3) with a mean average of 19.6mm zone of inhibition[fig 1]

www.earthjournals.org

Volume 4 Issue 2, 2014



Zone of inhibition of the control group of clotrimazole (B1, B2 and B3) with a mean diameter of 23.2mm [fig 2]

Table 1: Showing diameter of	zone of inhibition for	virgin coconut oil	and clotrimazole
at 24 hours:			

Group	A1	A2	A3	B1	B2	B3
1						
Inhibition	10,00	$21.5 \pm 0.5$	10.2 0 7	225105	24.1.0.9	$22.2 \pm 0.7$
Infilibition	18±0.8	21.3±0.3	19.5±0.7	23.3±0.3	24.1±0.8	$22.2\pm0.7$
zone	mm	mm	mm	mm	mm	mm
A	10 (			02 5		
Average	19.0 mm			23.5mm		

#### **DISCUSSION:**

Candida albicans is a harmless commensal yeast-like fungus in healthy humans, which can cause superficial as well as life threatening systemic infections in immunocompromised situations. C. albicans can colonize or infect virtually all body sites because of its high adaptability to different host niches by the activation of appropriate sets of genes in response to complex environmental signals. Oral candidiasis is common in the elderly, and in HIV infected patients. Secondary candidiasis also occurs in patients with long-term use of broad-spectrum antibiotics, corticosteroids, and immune-

www.earthjournals.org	Volume 4 Issue 2, 2014	29
www.earthjournals.org	Volume 4 Issue 2, 2014	29

suppressants.<sup>11, 12</sup> In these cases, antifungal agents are often prescribed. Recently Candida species have been shown increasing resistance to antifungals.

Coconut oil is edible oil prolific globally in tropical countries. The oil can be processed and extracted as an organic product which, is employed in the cosmetic industry for skin care to moisturize the skin, relieve dryness, flaking and prevent stretch marks. It is used for wounds, bruises, burns, rashes, eczema, and dermatitis. It also supports the natural chemical balance of the skin and provides protection from the damaging effects of ultraviolet radiation from the sun. Coconut, due to its contents of caprylic acid, which is fungicidal, is used in the treatment of fungal skin infections such as athlete's foot, thrush, ringworm and candidiasis.<sup>10, 13</sup>

Virgin coconut oil is derived from the milk obtained from fresh coconut, by processes like fermentation, centrifugal separation and enzyme actions. They grate the fresh coconut, extract the coconut milk, and then let the coconut milk stand in a covered container for about a day. After 24-36 hours, the oil naturally separates from the water producing crystal clear oil that retains the full scent and taste of coconuts. The active ingredients in Coconut Oil are Caprylic Acid, Capric Acid and Lauric Acid. These medium-chain saturated fatty acids are potent antimicrobial agents.<sup>14</sup> Lauric acid is the precursor of monolaurin (Pereira et al., 2004), which has been shown to modulate immune cell proliferation (Witcher et al., 1996) and possess antimicrobial activity (Bergsson et al., 2002). They kill bacteria, viruses and fungi by causing their cell walls to disintegrate. Several studies show how all the fatty acids demonstrate antifungal activity against C. albicans. Our present study shows virgin coconut oil as an effective alternative to Clotrimazole with the added advantage of being less costly, having no side effects, better acceptability by patients, easy availability and ease of application.

#### **CONCLUSION:**

The results of this study indicate that virgin coconut oil exerts an antifungal effect on Candida albicans. The findings as reported in the present study indicate a strong potential therapeutic value of virgin coconut oil against Candidial infections. Oral therapeutic products from coconut oil should be safe, effective and efficient alternative in antifungal therapy clinically. This will certainly increase the quality of life and decrease morbidity of the chronic fungal oral lesion that are seen associated with several debilitating diseases especially in the extremes of life.

#### **REFERENCES:**

- 1. Ramage G, Vandewalle K, Wickes BL, Lopez-Ribot JL. Characteristics of biofilm formation by Candida albicans.Rev Iberoam Micol 2001; 18: 163–170.
- 2. Sandven P. Epidemiology of candidemia. Rev Iberoam Micol 2000; 17: 73-81.
- 3. Haberland-Carrodeguas C, Allen CM, Beck FM et al (2002) Prevalence of fluconazole-resistant strains of Candida albicans in otherwise healthy outpatients. J Oral Pathol Med 31:99–105
- Friedman S, Richardson SE, Jacobs SE, O'Brien K: Systemic infection in extremely low birth weight infants: Short term morbidity and neuro developmental outcome. Ped Infect Disease J 19: 499–504, 2000

www.earthjournals.org

Volume 4 Issue 2, 2014

- Mackenzie DWR, Cauwenberg G, Van Cutsem J, Drouhet E, Dupont B: Mycoses in AIDS patients: An overview. In: H. Vanden Bossche (ed), Mycoses in AIDS Patients. New York, Plenum Press, 1990, p 27–33
- 6. Donlan RM. Biofilm formation: a clinically relevant microbiological process. Clin Infect Dis 2001; 33: 1387–1392.
- 7. Douglas LJ. Candida biofilms and their role in infection. Trends Microbiol 2003; 11: 30-36.
- 8. Kojic EM, Darouiche RO. Candida infections of medical devices. Clin Microbiol Rev 2004; 17: 255–267.
- Enig MG Lauric oils as antimicrobial agents: Theory of effect, scientific rationale and dietary application as adjunct nutrition support for HIV individuals, In: Nutrients and Food in AIDS. CRC Press, Boca Raton, 1999; 81-97.
- 10. Fife B The Healing Miracles of Coconut Oil. Piccadilly Books Ltd, Health wise publications, Colorado Springs, Co., 2000; 1-46.
- 11. Manohar V, Sirsi M, Ramananda Rao G: Yeasts in superficial mycoses . Incidence, isolation and characterization. Indian J Med Res 63: 261–265, 1975
- 12. Manohar V, Sirsi M, Ramananda Rao G: Yeasts in superficial mycosis II. Pathogenicity of Candida species to swiss albino mice. Indian J Med Res 64: 9–15, 1975
- 13. Bergsson G In vitro killing of Candida albicans by fatty acids and monoglycendes. Antimicrob Agents and Chemo. 2001; 45(11): 3209-3212.
- 14. Gudmundur Bergsson, et.al. (November 2001)."In Vitro killing of Candida albicans by fatty acids and monoglycerides".antimicrobial agents and chemotherapy volume 45:3209-3212, No.11 medicinal food volume 10:384-387, no.2.
- 15. Ogbulu,D.O.,et.al.(June 2007).In Vitro antimicrobial properties of coconut oil on Candida species in Ibadan,Nigeria.journal of medicinal food volume 10:384-387,no.2.